Read this package insert completely before use and comply with the instructions for use.

HealiAid® Collagen Wound Dressing

GRAPHICAL SYMBOLS



Single use only



Read instructions before use Sterilization by



irradiation Do not re-sterilize



Use by expiry date



Manufacturer



LOT

Do not use if package is damaged

Catalog number

Lot number



RONLY Caution: Federal law(USA) restricts this device to sale by or on the order of a licensed healthcare practitioner

Store at room temperature

Keep away from sunlight

Description

HealiAid® Collagen Wound Dressing is a white, porous, pliable and absorbable collagen wound dressing. It is fabricated by fibrous collagen matrix, in which the collagen is purified from bovine Achilles tendons. HealiAid® Collagen Wound Dressing is pliable and can be applied easily to clean wounds. HealiAid® Collagen Wound Dressing is supplied sterile. non-pyrogenic, and individually packaged. HealiAid® Collagen Wound Dressing in for single use only.

Indications

HealiAid® Collagen Wound Dressing is indicated for use in patients who have surgical wounds, donor site/grafts, podiatric wounds, wound dehiscence, traumatic wounds, abrasions, lacerations, partial thickness burns or skin tears. HealiAid® Collagen Wound Dressing can be applied to wounds with depth less than 0.3 cm.

Instructions for use

applying to the wound.

- · Apply HealiAid® Collagen Wound Dressing to cover wound site:
 - 1.Clean wounds with appropriate wound cleanser.
 - 2.HealiAid® Collagen Wound Dressing should be applied directly on the surface of the wound site. If the wound is dry (with drainage), HealiAid® Collagen Wound Dressing can be moistened before
 - 3. Cover the wound with appropriate cover dressing.
 - 4.Use HealiAid® Collagen Wound Dressing products until the wound is fully healed.
- · Removal of HealiAid® Collagen Wound Dressing:
 - 1.Remove the cover dressing and any remaining HealiAid® Collagen Wound Dressing carefully. In the case when HealiAid® Collagen Wound Dressing adheres to the wound site, rinse the wound site with normal saline before peeling off. Remove HealiAid® from the wound site carefully.
 - 2. The wound site should be cleaned with normal saline before replacing HealiAid® Collagen Wound Dressing. Dry the wound site before applying a new HealiAid®.
 - 3.Repeat replacing new HealiAid® Collagen Wound Dressing products to the wound dressing when needed.

Precautions

- HealiAid® Collagen Wound Dressing should not be used on infected
- Clean the wound site carefully before applying HealiAid® Collagen Wound Dressing.
- HealiAid® Collagen Wound Dressing is not intended to use in patients with abnormal coagulation.
- The safety of using HealiAid® Collagen Wound Dressing on pregnant women is not clinically evaluated. Surgeon should evaluate the application of HealiAid® Collagen Wound Dressing on pregnant women.
- Patients with severe allergy history or hypersensitive to bovine collagen are not recommended to use HealiAid®.
- HealiAid® Collagen Wound Dressing is contraindicated to patients undergoing desensitization injections to meat products, as these injections may contain bovine collagen.
- Check the expiration date of individual HealiAid® before use. Do not use any expired products.
- Do not use HealiAid® Collagen Wound Dressing when packaging is damaged, as the sterility of the product cannot be assured.
- HealiAid® Collagen Wound Dressing is restricted to single use only. Do not re-sterilized or re-used HealiAid® Collagen Wound Dressing.

Specifications

Specification
15 mm x 20 mm x 0.5 mm
20 mm x 30 mm x 0 5 mm
25 mm x 30 mm x 0.5 mm
30 mm x 40 mm x 0.5 mm
100 mm x 100 mm x 0.5 mm
20 mm x 40 mm x 3.0 mm
30 mm x 30 mm x 3 0 mm
40 mm x 50 mm x 3.0 mm
50 mm x 50 mm x 3 0 mm
100 mm x 100 mm x 3.0 mm
1 cc
2 cc
5 cc
10 cc
20 cc

Storage

HealiAid® Collagen Wound Dressing should be stored below 25°C (77°F) in a dry place. Do not freeze or expose HealiAid® to extreme conditions.

Warnings

HealiAid® Collagen Wound Dressing cannot be autoclaved because the collagen fibrous structure as well as the packaging will be damaged. HealiAid® Collagen Wound Dressing should not be re-sterilized. Residual portions of HealiAid® Collagen Wound Dressing should be discarded after use.

HealiAid® Collagen Wound Dressing contains bovine collagen and must not use in patients with a history of allergies to any bovine collagen products to prevent hypersensitivity. Hypersensitivity reactions reported when using bovine collagen products including erythema, swelling, induration, and/or urticaria.



MAXIGEN BIOTECH INC.

No.88, Keji 1st Rd., Guishan District, Taoyuan City 33383, Taiwan

TEL: +886-3-3287-222 FAX: +886-3-3287-333

F-MAIL : service@mbi-bio.com 21H2B0022